



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2019

Search for Light Dark Matter Interactions Enhanced by the Migdal Effect or Bremsstrahlung in XENON1T

Baudis, Laura ; Brown, Adam ; Galloway, Michelle ; Kish, A ; Manfredini, Alessandro ; Peres, R ;
Reichard, Shayne ; Volta, Giovanni ; Wulf, J ; et al

Abstract: Direct dark matter detection experiments based on a liquid xenon target are leading the search for darkmatter particles with masses above $5\text{GeV}/c^2$, but have limited sensitivity to lighter masses because of the small momentum transfer in dark matter-nucleus elastic scattering. However, there is an irreducible contribution from inelastic processes accompanying the elastic scattering, which leads to the excitation and ionization of the recoiling atom (the Migdal effect) or the emission of a bremsstrahlung photon. In this Letter, we report on a probe of low-mass dark matter with masses down to about $85\text{MeV}/c^2$ by looking for electronic recoils induced by the Migdal effect and bremsstrahlung using data from the XENON1T experiment. Besides the approach of detecting both scintillation and ionization signals, we exploit an approach that uses ionization signals only, which allows for a lower detection threshold. This analysis significantly enhances the sensitivity of XENON1T to light dark matter previously beyond its reach

DOI: <https://doi.org/10.1103/physrevlett.123.241803>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-178995>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution 4.0 International (CC BY 4.0) License.

Originally published at:

Baudis, Laura; Brown, Adam; Galloway, Michelle; Kish, A; Manfredini, Alessandro; Peres, R; Reichard, Shayne; Volta, Giovanni; Wulf, J; et al (2019). Search for Light Dark Matter Interactions Enhanced by the Migdal Effect or Bremsstrahlung in XENON1T. *Physical Review Letters*, 123(24):241803.

DOI: <https://doi.org/10.1103/physrevlett.123.241803>

Search for Light Dark Matter Interactions Enhanced by the Migdal Effect or Bremsstrahlung in XENON1T

E. Aprile,¹ J. Aalbers,² F. Agostini,³ M. Alfonsi,⁴ L. Althueser,⁵ F. D. Amaro,⁶ V. C. Antochi,² E. Angelino,⁷ F. Arneodo,⁸ D. Barge,² L. Baudis,⁹ B. Bauermeister,² L. Bellagamba,³ M. L. Benabderrahmane,⁸ T. Berger,¹⁰ P. A. Breur,¹¹ A. Brown,⁹ E. Brown,¹⁰ S. Bruenner,¹² G. Bruno,⁸ R. Budnik,¹³ C. Capelli,⁹ J. M. R. Cardoso,⁶ D. Cichon,¹² D. Coderre,¹⁴ A. P. Colijn,^{11,*} J. Conrad,² J. P. Cussonneau,¹⁵ M. P. Decowski,¹¹ P. de Perio,¹ A. Depoian,¹⁶ P. Di Gangi,³ A. Di Giovanni,⁸ S. Diglio,¹⁵ A. Elykov,¹⁴ G. Eurin,¹² J. Fei,¹⁷ A. D. Ferella,² A. Fieguth,⁵ W. Fulgione,^{18,7} P. Gaemers,¹¹ A. Gallo Rosso,¹⁸ M. Galloway,⁹ F. Gao,¹ M. Garbini,³ L. Grandi,¹⁹ Z. Greene,¹ C. Hasterok,¹² C. Hils,⁴ E. Hogenbirk,¹¹ J. Howlett,¹ M. Iacovacci,²⁰ R. Itay,¹³ F. Joerg,¹² S. Kazama,^{21,†} A. Kish,⁹ M. Kobayashi,¹ G. Koltman,¹³ A. Kopec,¹⁶ H. Landsman,¹³ R. F. Lang,¹⁶ L. Levinson,¹³ Q. Lin,^{1,‡} S. Lindemann,¹⁴ M. Lindner,¹² F. Lombardi,^{6,17} J. A. M. Lopes,^{6,8} E. López Fune,²² C. Macolino,²³ J. Mahlstedt,² M. Manenti,⁸ A. Manfredini,^{9,13} F. Marignetti,²⁰ T. Marrodán Undagoitia,¹² J. Masbou,¹⁵ S. Mastroianni,²⁰ M. Messina,^{18,8} K. Micheneau,¹⁵ K. Miller,¹⁹ A. Molinaro,¹⁸ K. Morā,² Y. Mosbacher,¹³ M. Murra,⁵ J. Naganoma,^{18,24} K. Ni,¹⁷ U. Oberlack,⁴ K. Odgers,¹⁰ J. Palacio,¹⁵ B. Pelssers,² R. Peres,⁹ J. Pienaar,¹⁹ V. Pizzella,¹² G. Plante,¹ R. Podvianiuk,¹⁸ J. Qin,¹⁶ H. Qiu,¹³ D. Ramírez García,¹⁴ S. Reichard,⁹ B. Riedel,¹⁹ A. Rocchetti,¹⁴ N. Rupp,¹² J. M. F. dos Santos,⁶ G. Sartorelli,³ N. Šarčević,¹⁴ M. Scheibelhut,⁴ S. Schindler,⁴ J. Schreiner,¹² D. Schulte,⁵ M. Schumann,¹⁴ L. Scotto Lavina,²² M. Selvi,³ P. Shagin,²⁴ E. Shockley,¹⁹ M. Silva,⁶ H. Simgen,¹² C. Therreau,¹⁵ D. Thers,¹⁵ F. Toschi,¹⁴ G. Trincherro,⁷ C. Tunnell,²⁴ N. Upole,¹⁹ M. Vargas,⁵ G. Volta,⁹ O. Wack,¹² H. Wang,²⁵ Y. Wei,¹⁷ C. Weinheimer,⁵ D. Wenz,⁴ C. Wittweg,⁵ J. Wulf,⁹ J. Ye,¹⁷ Y. Zhang,¹ T. Zhu,¹ and J. P. Zopounidis²²

(XENON Collaboration)^{||}

¹Physics Department, Columbia University, New York, New York 10027, USA

²Oskar Klein Centre, Department of Physics, Stockholm University, AlbaNova, Stockholm SE-10691, Sweden

³Department of Physics and Astronomy, University of Bologna and INFN-Bologna, 40126 Bologna, Italy

⁴Institut für Physik and Exzellenzcluster PRISMA, Johannes Gutenberg-Universität Mainz, 55099 Mainz, Germany

⁵Institut für Kernphysik, Westfälische Wilhelms-Universität Münster, 48149 Münster, Germany

⁶LIBPhys, Department of Physics, University of Coimbra, 3004-516 Coimbra, Portugal

⁷INAF-Astrophysical Observatory of Torino, Department of Physics, University of Torino and INFN-Torino, 10125 Torino, Italy

⁸New York University Abu Dhabi, P.O. Box 129188, Abu Dhabi, United Arab Emirates

⁹Physik-Institut, University of Zurich, 8057 Zurich, Switzerland

¹⁰Department of Physics, Applied Physics and Astronomy, Rensselaer Polytechnic Institute, Troy, New York 12180, USA

¹¹Nikhef and the University of Amsterdam, Science Park, 1098XG Amsterdam, Netherlands

¹²Max-Planck-Institut für Kernphysik, 69117 Heidelberg, Germany

¹³Department of Particle Physics and Astrophysics, Weizmann Institute of Science, Rehovot 7610001, Israel

¹⁴Physikalisches Institut, Universität Freiburg, 79104 Freiburg, Germany

¹⁵SUBATECH, IMT Atlantique, CNRS/IN2P3, Université de Nantes, Nantes 44307, France

¹⁶Department of Physics and Astronomy, Purdue University, West Lafayette, Indiana 47907, USA

¹⁷Department of Physics, University of California, San Diego, California 92093, USA

¹⁸INFN-Laboratori Nazionali del Gran Sasso and Gran Sasso Science Institute, 67100 L'Aquila, Italy

¹⁹Department of Physics and Kavli Institute for Cosmological Physics, University of Chicago, Chicago, Illinois 60637, USA

²⁰Department of Physics "Ettore Pancini," University of Napoli and INFN-Napoli, 80126 Napoli, Italy

²¹Kobayashi-Maskawa Institute for the Origin of Particles and the Universe, Nagoya University, Furo-cho, Chikusa-ku, Nagoya, Aichi 464-8602, Japan

²²LPNHE, Université Pierre et Marie Curie, Université Paris Diderot, CNRS/IN2P3, Paris 75252, France

²³*LAL, Université Paris-Sud, CNRS/IN2P3, Université Paris-Saclay, F-91405 Orsay, France*²⁴*Department of Physics and Astronomy, Rice University, Houston, Texas 77005, USA*²⁵*Physics and Astronomy Department, University of California, Los Angeles, California 90095, USA*

(Received 31 July 2019; revised manuscript received 15 October 2019; published 13 December 2019)

Direct dark matter detection experiments based on a liquid xenon target are leading the search for dark matter particles with masses above $\sim 5 \text{ GeV}/c^2$, but have limited sensitivity to lighter masses because of the small momentum transfer in dark matter-nucleus elastic scattering. However, there is an irreducible contribution from inelastic processes accompanying the elastic scattering, which leads to the excitation and ionization of the recoiling atom (the Migdal effect) or the emission of a bremsstrahlung photon. In this Letter, we report on a probe of low-mass dark matter with masses down to about $85 \text{ MeV}/c^2$ by looking for electronic recoils induced by the Migdal effect and bremsstrahlung using data from the XENON1T experiment. Besides the approach of detecting both scintillation and ionization signals, we exploit an approach that uses ionization signals only, which allows for a lower detection threshold. This analysis significantly enhances the sensitivity of XENON1T to light dark matter previously beyond its reach.

DOI: [10.1103/PhysRevLett.123.241803](https://doi.org/10.1103/PhysRevLett.123.241803)

The existence of dark matter (DM) is supported by various astronomical and cosmological observations [1–3], but its nature remains unknown. The most promising DM candidate is the so-called weakly interacting massive particle (WIMP) [4], which explains the current abundance of dark matter as a thermal relic of the big bang [5]. In the past three decades, numerous terrestrial experiments have been built to detect the faint interactions between WIMPs and ordinary matter. Among them, experiments using dual-phase (liquid-gas) xenon time projection chambers (TPCs) [6–8] are leading the search for WIMPs with masses from a few GeV/c^2 to TeV/c^2 . The mass of the WIMP is expected to be larger than about $2 \text{ GeV}/c^2$ from the Lee-Weinberg limit [5] assuming a weak scale interaction. On the other hand, DM in the sub- GeV/c^2 mass range has more recently been proposed in several models [9]. In this Letter, we report on a probe of light DM-nucleon elastic interactions by looking for electronic recoils (ERs) in XENON1T, induced by secondary radiation, bremsstrahlung (BREM) [10] and the Migdal effect [11,12], that can accompany a nuclear recoil (NR). ER signals induced by the Migdal effect and BREM can go well below 1 keV, where the detection efficiency for the scintillation signal is low. Therefore, in addition to the analysis utilizing both ionization and scintillation signals, we performed an analysis using the ionization signal only, which improves the detection efficiency for sub-keV ER events. We present results from a probe of light DM (LDM) with masses as low as $85 \text{ MeV}/c^2$.

The XENON1T direct dark matter detection experiment [13] uses a dual-phase TPC containing 2 tonnes of ultrapure liquid xenon (LXe) as the active target material. It is located at the INFN Laboratori Nazionali del Gran Sasso (LNGS) in Italy, which has an average rock overburden of 3600 m water equivalent. The prompt primary scintillation (S1) and secondary electroluminescence of ionized electrons (S2) signals are detected by top and bottom arrays of 248 Hamamatsu R11410-21 3" photomultiplier tubes (PMTs) [14,15]. They are used to reconstruct the deposited energy and the event interaction position in three dimensions, which allows for fiducialization of the active volume [16,17]. The XENON1T experiment has published WIMP search results by looking for NRs from WIMP-nucleus elastic scattering using data from a one-tonne-year exposure, achieving the lowest ER background in a DM search experiment [8]. The excellent sensitivity of LXe experiments to heavy WIMPs comes from the heavy xenon nucleus which gives a coherent enhancement of the interaction cross section and from the large NR energy. The sensitivity to sub- GeV/c^2 LDM, on the other hand, decreases rapidly with lowering DM mass since detectable scintillation and ionization signals produced by these NRs become too small. The energy threshold (defined here as the energy at which the efficiency is 10%) in a LXe TPC is mainly limited by the amount of detectable S1 signals. A significant fraction of deposited NR energy is transferred into heat due to the Lindhard quenching effect [18]. Thus the detection efficiency for these NRs becomes extremely low, with less than 10% for NRs below 3.5 keV in XENON1T [8]. It is challenging to detect the NR signals from LDM interactions.

Unlike NRs, ERs lose negligible energy as heat because recoil electrons have small masses compared with xenon nuclei. This leads to a lower energy threshold for ER signals. Probing the ER signals induced by the Migdal

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/). Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

effect (MIGD) and BREM enables a significant boost of XENON1T's sensitivity to LDMs, thanks to the lowered threshold.

When a particle elastically scatters off a xenon nucleus, the nucleus undergoes a sudden momentum change with respect to the orbital atomic electrons, resulting in the polarization of the recoiling atom and a kinematic boost of the electrons. The depolarization process can lead to BREM emission [10], and the kinematic boost of atomic electrons can result in ionization and/or excitation of the atom, which eventually causes secondary radiation, known as the Migdal effect [11,12].

The differential rate of BREM emission with photon energy E_{ER} is given by

$$\frac{d^2 R}{dE_{\text{ER}} dv} \propto \frac{|f(E_{\text{ER}})|^2}{E_{\text{ER}}} \sqrt{1 - \frac{2E_{\text{ER}}}{\mu_N v^2} \left(1 - \frac{E_{\text{ER}}}{\mu_N v^2}\right)}, \quad (1)$$

where v , μ_N , and $f(E_{\text{ER}})$ are the velocity of DM, the reduced mass of the xenon nucleus and DM, and the atomic scattering factor, respectively [10].

The differential rate of the MIGD process giving a NR of energy E_{NR} accompanied by an ER of energy E_{ER} is given by

$$\frac{dR}{dE_{\text{ER}}} \simeq \int dE_{\text{NR}} dv \frac{d^2 R}{dE_{\text{NR}} dv} \times \frac{1}{2\pi} \sum_{n,l} \frac{d}{dE_{\text{ER}}} p_{q_e}^c(n, l \rightarrow E_{\text{ER}} - E_{n,l}), \quad (2)$$

where $p_{q_e}^c$ is the probability for an atomic electron, with quantum numbers (n, l) and binding energy $E_{n,l}$, to be ionized and receive a kinetic energy $E_{\text{ER}} - E_{n,l}$ [12]. $p_{q_e}^c$ is related to q_e , which is the momentum of each electron in the rest frame of the nucleus after the scattering. The shell vacancy is immediately refilled, and an x ray or an Auger electron with energy $E_{n,l}$ is emitted. $E_{n,l}$ is measured simultaneously with the energy deposited by the ionized electron, since the typical timescale of the deexcitation process is $\mathcal{O}(10)$ fs. Atomic electrons can also undergo excitation instead of ionization, in which case an x ray is emitted during deexcitation [12]. Excitation, however, is subdominant compared to the ionization process, and thus is not considered in this analysis. Only the contributions from the ionization of M -shell ($n = 3$) and N -shell ($n = 4$) electrons are considered in this work, as inner electrons ($n \leq 2$) are too strongly bound to the nucleus to contribute significantly. The contribution from the ionization of valence electrons ($n = 5$) is neglected because it is subdominant in the region of interest compared to the ones from M - and N -shell electrons, and the calculation of it has large uncertainty since the assumption of isolated atoms is used for LXe [12]. An illustration of MIGD and BREM is given in Fig. 1. The radiation from MIGD is typically 3–4




FIG. 1. Illustration of the ER signal production from BREM (green) and MIGD processes (pink) after elastic scattering between DM (χ) and a xenon nucleus. The electrons illustrated in pink represent those involved in ionization, deexcitation, and Auger electron emission during a MIGD process.

orders of magnitude more likely to occur than BREM. Although only a very small fraction (about 3×10^{-8} and 8×10^{-6} for DM masses of 0.1 and 1.0 GeV/c^2 , respectively) of NRs accompanies MIGD radiations, the larger energy and ER nature make them easier to be detected than the pure NRs.

The data used in previous analyses [8] consist of two science runs with a live time of 32.1 days (SR0) and 246.7 days (SR1), respectively. The two runs were taken under slightly different detector conditions. To maximize the amount of data acquired under stable detector conditions, we decided to use SR1 only. The same event selection, fiducial mass, correction, and background models as described in Ref. [8] are used for the SR1 data, which we refer to as the S1-S2 data in later text. The exposure of the S1-S2 data is about 320 tonne-days. The interpretation of such S1-S2 analysis is based on the corrected S1 (cS1) signal and the corrected S2 signal from the PMTs at the bottom of the TPC (cS2_b).

The region of interest in the S1-S2 data is from 3 to 70 photoelectrons (PEs) in cS1, which corresponds to median ER energies from 1.4 to 10.6 keV in the 1.3-tonne fiducial volume of XENON1T. The lower value is dictated by the requirement of the threefold PMT coincidence for defining a valid S1 signal [16]. A detailed signal response model [17] is used to derive the influence of various detector features, including the requirement of the threefold PMT coincidence, on the reconstructed signals. The effective exposure, which is defined as exposure times detection efficiency, and its uncertainty as a function of deposited ER energy for the S1-S2 data are shown in Fig. 2, with the signal spectra from MIGD and BREM induced by 0.1 and 1 GeV/c^2 DM masses overlaid. The (cS2_b, cS1) distribution of S1-S2 data are shown in Fig. 3. The rise of the event rate at around 0.85 keV for DM mass of 1.0 GeV/c^2 is contributed by the ionization of M -shell electrons [10,12].




FIG. 2. Median effective exposures of ER signals after event selections as a function of recoil energy for the $S1$ - $S2$ data (black line) and $S2$ -only data (red line). The 68% credible regions of the effective exposures are also shown as the shaded regions. The expected event rate of DM-nucleus scattering from MIGD (BREM) for DM masses of 0.1 and 1.0 GeV/c^2 are overlaid as well, in magenta (green) dashed and solid lines, respectively, assuming a spin-independent DM-nucleon interaction cross section of 10^{-35} cm^2 .

In our signal models, deposited energy below 1 keV, at which the median detection efficiency in 1.3-tonne fiducial volume is 10%, from MIGD and BREM is neglected for the $S1$ - $S2$ data in the following analysis. There are only two sub-keV measurements of ionization yield for ER in LXe [19,20].

The $S1$ - $S2$ data selections [16] provide excellent rejection of noise and backgrounds, and are characterized as well by the well-established background models [17] and a fully blind analysis [8]. However, they also limit the detection efficiency of $\mathcal{O}(1)$ keV energy depositions. We therefore consider also the events with no specific requirement on $S1$ ($S2$ -only data) in this work. Although the reduction of available information in the $S2$ -only data implies less background discrimination, the increased detection efficiency in the < 1 keV ER energy region, shown in Fig. 2, enables a more sensitive search for LDM-nucleus interactions through MIGD and BREM. The interpretation of such $S2$ -only data is based on the uncorrected $S2$ signal, combining both signals from top and bottom PMT arrays.

We analyze the $S2$ -only data as in Ref. [21], using the LDM signal models appropriate for MIGD and BREM. As detailed in Ref. [21], 30% of the data were used for choosing regions of interest (ROIs) in $S2$ and event selections. A different $S2$ ROI is chosen for each dark matter model and mass to maximize the signal-to-noise ratio, based on the training data. The event selections used for this work are the same as in Ref. [21], and mainly based on the width of each $S2$ waveform, reconstructed radius, and PMT hit pattern of the $S2$. Figure 4 shows the observed $S2$ spectra for the $S2$ -only data, along with the expected




FIG. 3. Comparison of $cS1$ and $cS2_b$ spectra between the $S1$ - $S2$ data and the signal response model [17]. In upper panel (I), the distribution of the $S1$ - $S2$ data in $(cS2_b, cS1)$ space is shown as light blue dots, along with the best-fit ER background model (black shaded region). The contours containing 90% of the expected signals from MIGD for 0.3, 0.5, and 1 GeV/c^2 DM are shown in red dotted, dashed, and solid lines, respectively. Gray lines show isoenergy contours in ER energy. The events having lower $cS2_b$ than what we expect for ER are mostly surface backgrounds [8], which have minimal impact to the results of this study. The lower panel (II) shows the projected $cS1$ distribution of the $S1$ - $S2$ data, where $cS2_b$ is within the 2σ contour of ER model shown in (I). For comparison, the 68% credible region of $cS1$ distribution from ER background model (blue shadow) is shown, which is mainly attributed to the systematic uncertainties of the model. The $cS1$ distributions of the expected signals from MIGD for 0.3, 0.5, and 1 GeV/c^2 DM with assumed spin-independent DM-nucleon cross sections of 2×10^{-28} , 10^{-36} , and 10^{-38} cm^2 , respectively, are shown as well. The vertical dashed lines indicate the region of interest (3–70 PE). The inset, panel (III), shows the $cS2_b$ distribution, with $cS1$ in (3,10) PE, compared with the 68% credible region of the $cS2_b$ spectra from the ER background model (blue shadow).

DM signal distributions by MIGD with masses of 0.1, 0.5, and 1.0 GeV/c^2 , respectively. The $S2$ ROIs for these three DM models shown in Fig. 4 are indicated by the colored arrows. Conservative estimates of the background from ^{214}Pb -induced β decays, solar-neutrino-induced NRs, and surface backgrounds from the cathode electrode are used in the inference [21]. The background model is shown in Fig. 4 as a shaded gray region.




FIG. 4. Observed $S2$ spectra for the $S2$ -only data after the optimized selection described in Ref. [21]. The expected spectra of ER signals induced by MIGD for DM with mass of 0.1, 0.5, and 1.0 GeV/c^2 are shown in green, blue, and red solid lines, respectively, assuming the spin-independent DM-nucleon interaction cross section of 10^{-33} cm^2 for 0.1 GeV/c^2 DM and of 10^{-35} cm^2 for 0.5 and 1.0 GeV/c^2 DM. The gray shaded region shows the conservative background model used in analysis of $S2$ -only data. The arrows indicate the $S2$ ROIs that are later used in inference for the three DM signals mentioned above. The $S2$ threshold used for the $S2$ -only data is denoted in the dashed black line.

The detector response to ERs from MIGD and BREM in ($cS2_b$, $cS1$) space (for the $S1$ - $S2$ data) and in reconstructed number of electrons (for the $S2$ -only data) is derived using the signal response model described in Ref. [17]. Note that the ionization yield used for the $S2$ -only data is more conservative than the noble element simulation technique (NEST) v2 model [22]. Figure 3 shows the comparison between the expectation from our signal response model and the $S1$ - $S2$ data, as well as the ($cS2_b$, $cS1$) distribution of ERs from MIGD. Signal contours for different DM masses are similar since the energy spectra from MIGD and BREM are not sensitive to incident dark matter velocity as long as it is kinematically allowed. We have ignored the contribution of NRs in the signal model of MIGD and BREM, since it is small compared with ERs from MIGD and BREM in this analysis and there is no measurement of scintillation and ionization yields in LXe for simultaneous ER and NR energy depositions. We use the inference only for DM mass below 2 GeV/c^2 , above which the contribution of a NR in the signal rate becomes comparable with or exceeds the signal model uncertainty.

The $S1$ - $S2$ data are interpreted using an unbinned profile likelihood ratio as the test statistic, as detailed in Ref. [17]. The unbinned profile likelihood is calculated using background models defined in $cS2_b$, $cS1$, and spatial coordinates. The uncertainties from the scintillation and ionization yields of ER backgrounds, along with the uncertainties in the estimated rates of each background component, are taken into account in the inference [17]. The inference procedure for the $S2$ -only data is detailed in

Ref. [21], which is based on simple Poisson statistics using the number of events in the $S2$ ROI. The event rates of spin-independent (SI) and -dependent (SD) DM-nucleon elastic scattering are calculated following the approaches described in Refs. [8,23] and Ref. [24], respectively.

The results are also interpreted in a scenario where LDM interacts with the nucleon through a scalar force mediator ϕ with equal effective couplings to the proton and neutron as in the SI DM-nucleon elastic scattering. In this scenario, the differential event rates are corrected by $m_\phi^4/(m_\phi^2 + q^2/c^2)^2$ [25,26], where $q = \sqrt{2m_N E_R}$ and m_N are the momentum transfer and the nuclear mass, respectively. We take the light mediator (LM) regime where the momentum transfer is much larger than m_ϕ and thus the interaction cross section scales with m_ϕ^4 . In this regime, the contribution of NRs is largely suppressed compared with SI DM-nucleon elastic scattering due to the long-range nature of the interaction. Therefore, the results are interpreted for DM mass up to 5 GeV/c^2 for SI-LM DM-nucleon elastic scattering.

In addition, we also take into account the fact that a DM particle may be stopped or scatter multiple times when passing through Earth's atmosphere, mantle, and core before reaching the detector (Earth-shielding effect) [27–29]. If the DM-matter interaction is sufficiently strong, the sensitivity for detecting such DM particles in terrestrial detectors, especially in an underground laboratory, can be reduced or even lost totally. Following Ref. [30], the VERNE code [31] is used to calculate the Earth-shielding effect for SI DM-nucleon interaction. A modification of the VERNE code based on the methodology in Ref. [32] is applied for the calculations of SD and SD-LM DM-nucleon interactions. To account for the Earth-shielding effect for SD DM-nucleon interaction, ^{14}N in the atmosphere and ^{29}Si in Earth's mantle and core are considered, and their spin expectation values, $\langle S_n \rangle$ and $\langle S_p \rangle$, are taken from Ref. [33]. Both the lower and upper boundaries of excluded parameter space are reported in this work. The lower boundaries are conventionally referred to as upper limits in later context, and are the primary interest of this work. The upper boundaries are dominated by the overburden configuration of the Gran Sasso laboratory which hosts the detector.

No significant excess is observed above the background expectation in the search using the $S1$ - $S2$ data. Figure 5 shows the 90% confidence level (C.L.) limits [34] on the SI and SD (proton-only and neutron-only cases) DM-nucleon interaction cross section using signal models from MIGD and BREM with masses from about 85 MeV/c^2 to 2 GeV/c^2 , and Fig. 6 shows the 90% C.L. limits [34] on the SI-LM DM-nucleon interaction cross section with masses from about 100 MeV/c^2 to 5 GeV/c^2 . The sensitivity contours for the results derived using $S2$ -only data are not shown because of the conservativeness of the background model. The upper limits derived using the $S1$ - $S2$




FIG. 5. Limits on the SI (upper panel), SD proton-only (middle panel), and SD neutron-only (lower panel) DM-nucleon interaction cross sections at 90% C.L. using signal models from MIGD and BREM in the XENON1T experiment with the S1-S2 data (blue contours and lines) and S2-only data (black contours and lines). The solid and dashed (dotted) lines represent the lower boundaries (also referred to as upper limits) and MIGD (BREM) upper boundaries of the excluded parameter regions. Green and yellow shaded regions give the 1σ and 2σ sensitivity contours for upper limits derived using the S1-S2 data, respectively. The upper limits on the SI DM-nucleon interaction cross sections from LUX [35], EDELWEISS [30], CDEX [36], CRESST-III [37], NEWS-G [38], CDMSlite-II [39], and DarkSide-50 [40], and upper limits on the SD DM-nucleon interaction cross sections from CRESST [37,41] and CDMSlite [42] are also shown. Note that the limits derived using the S1-S2 and S2-only data are inferred using unbinned profile likelihood method [16] and simple Poisson statistics with the optimized event selection [21], respectively. The sensitivity contours for the S2-only data are not given since the background models used in the S2-only data are conservative [21].

data deviate from the median sensitivity by about 1σ – 2σ due to the underfluctuation of the ER background in the low energy region. As described in Ref. [21], the jumps in the S2-only limits are originating from the changes in the observed number of events due to the mass-dependent S2 ROIs. The results, by searching for ER signals induced by MIGD, give the best lower exclusion boundaries on SI, SD




FIG. 6. Limits on the SI-LM DM-nucleon interaction cross sections at 90% C.L. using signal models from MIGD and BREM in the XENON1T experiment with the S1-S2 data (blue contours and lines) and S2-only data (black contours and lines). The description is the same as in Fig. 5. The upper limits on the SI DM-nucleon interaction cross sections from LUX [35] and XENON1T S2-only (elastic NR results) [21] are also shown.

proton-only, SD neutron-only, and SI-LM DM-nucleon interaction cross section for mass below about 1.8, 2.0, 2.0, and 4.0 GeV/c^2 , respectively, as compared to previous experiments [30,35–42]. The upper limits derived from the S1-S2 data become comparable with those from the S2-only data at $\sim \text{GeV}/c^2$ since the efficiency of the S1-S2 data to DM signals with mass of $\sim \text{GeV}/c^2$ becomes sufficiently high. However, the upper limits derived from the S1-S2 data do not provide significantly better constraints than those from the S2-only data for DM masses larger than 1 GeV/c^2 , because both data are dominated by the ER background, which is very similar to the expected DM signal.

In summary, we performed a search for LDM by probing ER signals induced by MIGD and BREM, using data from the XENON1T experiment. These new detection channels significantly enhance the sensitivity of LXe experiments to masses unreachable in the standard NR searches. We set the most stringent upper limits on the SI and SD DM-nucleon interaction cross sections for masses below 1.8 and 2 GeV/c^2 , respectively. Together with the standard NR search [8], XENON1T results have reached unprecedented sensitivities to both low-mass (sub- GeV/c^2) and high-mass (GeV/c^2 – TeV/c^2) DM. With the upgrade to XENONnT, we expect to further improve the sensitivity to DM with masses ranging from about 85 MeV/c^2 to beyond a TeV/c^2 .

The authors would like to thank Masahiro Ibe and Yutaro Shoji for helpful discussions on MIGD and for providing us with the code for calculating the rate of MIGD radiation in xenon. We would like to thank Bradley Kavanagh for helpful discussion on the Earth-shielding effect. We gratefully acknowledge support from the National Science Foundation, Swiss National Science Foundation, German Ministry for Education and Research, Max Planck Gesellschaft, Deutsche Forschungsgemeinschaft,

Netherlands Organisation for Scientific Research (NWO), Netherlands eScience Center (NLeSC) with the support of the SURF Cooperative, Weizmann Institute of Science, Israeli Centers Of Research Excellence (I-CORE), Pazy-Vatat, Fundacao para a Ciencia e a Tecnologia, Region des Pays de la Loire, Knut and Alice Wallenberg Foundation, Kavli Foundation, and Istituto Nazionale di Fisica Nucleare. This project has received funding or support from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie Grant Agreements No. 690575 and No. 674896, respectively. Data processing is performed using infrastructures from the Open Science Grid and European Grid Initiative. We are grateful to Laboratori Nazionali del Gran Sasso for hosting and supporting the XENON project.

*Also at Institute for Subatomic Physics, Utrecht University, 3584 CC Utrecht, Netherlands.

†kazama@isee.nagoya-u.ac.jp

‡ql2265@vip.163.com

§Also at Coimbra Polytechnic—ISEC, 3030-199 Coimbra, Portugal.

||xenon@lngs.infn.it

- [1] D. Clowe, A. Gonzalez, and M. Markevitch, *Astrophys. J.* **604**, 596 (2004).
- [2] V. C. Rubin, W. K. Ford, Jr., and N. Thonnard, *Astrophys. J.* **238**, 471 (1980).
- [3] N. Aghanim *et al.* (Planck Collaboration), *arXiv:1807.06209*.
- [4] G. Jungman, M. Kamionkowski, and K. Griest, *Phys. Rep.* **267**, 195 (1996).
- [5] B. W. Lee and S. Weinberg, *Phys. Rev. Lett.* **39**, 165 (1977).
- [6] D. S. Akerib *et al.* (LUX Collaboration), *Phys. Rev. Lett.* **116**, 161302 (2016).
- [7] X. Cui *et al.* (PandaX-II Collaboration), *Phys. Rev. Lett.* **119**, 181302 (2017).
- [8] E. Aprile *et al.* (XENON Collaboration), *Phys. Rev. Lett.* **121**, 111302 (2018).
- [9] M. Battaglieri *et al.*, *arXiv:1707.04591*.
- [10] C. Kouvaris and J. Pradler, *Phys. Rev. Lett.* **118**, 031803 (2017).
- [11] A. B. Migdal, *J. Phys. (USSR)* **4**, 449 (1941).
- [12] M. Ibe, W. Nakano, Y. Shoji, and K. Suzuki, *J. High Energy Phys.* **03** (2018) 194.
- [13] E. Aprile *et al.* (XENON Collaboration), *Eur. Phys. J. C* **77**, 881 (2017).
- [14] E. Aprile *et al.* (XENON Collaboration), *Eur. Phys. J. C* **75**, 546 (2015).
- [15] E. Aprile *et al.* (XENON Collaboration), *J. Instrum.* **12**, P01024 (2017).
- [16] E. Aprile *et al.* (XENON Collaboration), *Phys. Rev. D* **100**, 052014 (2019).
- [17] E. Aprile *et al.* (XENON Collaboration), *Phys. Rev. D* **99**, 112009 (2019).
- [18] J. Lindhard, V. Nielsen, M. Scharff, and P. Thomsen, *Kgl. Danske Videnskab., Selskab. Mat. Fys. Medd.* **33** (1963).
- [19] D. S. Akerib *et al.* (LUX Collaboration), *Phys. Rev. D* **96**, 112011 (2017).
- [20] E. Boulton, E. Bernard, N. Destefano, B. Edwards, M. Gai, S. Hertel, M. Horn, N. Larsen, B. Tennyson, C. Wahl *et al.*, *J. Instrum.* **12**, P08004 (2017).
- [21] E. Aprile *et al.* (XENON Collaboration), *arXiv:1907.11485*.
- [22] M. Szydagis *et al.*, Noble element simulation technique v2.0, <https://doi.org/10.5281/zenodo.1314669> (2018).
- [23] J. Lewin and P. Smith, *Astropart. Phys.* **6**, 87 (1996).
- [24] E. Aprile *et al.* (XENON Collaboration), *Phys. Rev. Lett.* **122**, 141301 (2019).
- [25] E. Del Nobile, M. Kaplinghat, and H. B. Yu, *J. Cosmol. Astropart. Phys.* **10** (2015) 055.
- [26] X. Ren *et al.* (PandaX-II Collaboration), *Phys. Rev. Lett.* **121**, 021304 (2018).
- [27] T. Emken and C. Kouvaris, *Phys. Rev. D* **97**, 115047 (2018).
- [28] M. S. Mahdawi and G. R. Farrar, *arXiv:1712.01170*.
- [29] T. Emken and C. Kouvaris, *J. Cosmol. Astropart. Phys.* **10** (2017) 031.
- [30] E. Armengaud *et al.* (EDELWEISS Collaboration), *Phys. Rev. D* **99**, 082003 (2019).
- [31] B. J. Kavanagh, bradkav/verne: Release, <https://doi.org/10.5281/zenodo.1116305> (2017).
- [32] B. J. Kavanagh, *Phys. Rev. D* **97**, 123013 (2018).
- [33] D. Hooper and S. D. McDermott, *Phys. Rev. D* **97**, 115006 (2018).
- [34] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.123.241803> for data points of limits.
- [35] D. S. Akerib *et al.* (LUX Collaboration), *Phys. Rev. Lett.* **122**, 131301 (2019).
- [36] Z. Z. Liu *et al.* (CDEX Collaboration), *Phys. Rev. Lett.* **123**, 161301 (2019).
- [37] A. Abdelhameed *et al.* (CRESST Collaboration), *arXiv:1904.00498* [Phys. Rev. D (to be published)].
- [38] Q. Arnaud *et al.* (NEWS-G Collaboration), *Astropart. Phys.* **97**, 54 (2018).
- [39] R. Agnese *et al.* (SuperCDMS Collaboration), *Phys. Rev. Lett.* **116**, 071301 (2016).
- [40] P. Agnes *et al.* (DarkSide Collaboration), *Phys. Rev. Lett.* **121**, 081307 (2018).
- [41] A. Abdelhameed *et al.* (CRESST Collaboration), *Eur. Phys. J. C* **79**, 630 (2019).
- [42] R. Agnese *et al.* (CDMS Collaboration), *Phys. Rev. D* **97**, 022002 (2018).